

For Use in
Swine Feeds Only

Tylan[®] 40

Sulfa-G[™]

Tylosin Phosphate and Sulfamethazine
Elliptical Pellets

*

Net Weight **50 lbs**
(22.68 kg)

Type A Medicated Article

Do not feed undiluted.

Equivalent to 40 g tylosin per pound and 40 g sulfamethazine per pound.

For maintaining weight gains and feed efficiency in the presence of atrophic rhinitis; lowering the incidence and severity of *Bordetella bronchiseptica* rhinitis; prevention of swine dysentery (bloody scours) caused by *Serpulina hyodysenteriae*; control of swine pneumonias caused by bacterial pathogens (*Pasteurella multocida* and/or *Actinomyces pyogenes*).

Important: Must Be Thoroughly Mixed in Feeds Before Use.

Manufactured for:

Elanco Animal Health • A Division of Eli Lilly and Company
Indianapolis, IN 46285, USA

Questions or Comments Call 1-800-428-4441

*Elanco®, Tylan® 40 Sulfa-G™, and the diagonal color bar are trademarks of Eli Lilly and Company
BG 1305 AMB (L-I-Mar-01)

For Use in
Swine Feeds Only

Tylan 40[®] Sulfa-G[™]

Tylosin Phosphate and Sulfamethazine Elliptical Pellets

Net Weight **50 lbs**
(22.68 kg)

Type A Medicated Article

Do not feed undiluted.

Active Drug Ingredients: Tylosin (as tylosin phosphate)..... 40 g per lb
Sulfamethazine 8.8% (40 g per lb)

Ingredients: Roughage products, calcium carbonate, and mineral oil.

Important: Must Be Thoroughly Mixed in Feeds Before Use.

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

Warning: Tylan 40 Sulfa-G may be irritating to unprotected skin and eyes. When mixing and handling Tylan 40 Sulfa-G use protective clothing and impervious gloves. In case of accidental eye exposure, flush eyes with plenty of water. Exposed skin should be washed with plenty of soap and water. Remove and wash contaminated clothing. Seek medical attention if irritation becomes severe or persists. The material safety data sheet (MSDS) contains more detailed occupational safety information. To report adverse effects in users or obtain a copy of the MSDS, call 1-800-428-4441.

For maintaining weight gains and feed efficiency in the presence of atrophic rhinitis; lowering the incidence and severity of *Bordetella bronchiseptica rhinitis*; prevention of swine dysentery (bloody scours) caused by *Serpulina hyodysenteriae*; control of swine pneumonias caused by bacterial pathogens (*Pasteurella multocida* and/or *Actinomyces pyogenes*).

Mixing and Feeding Directions: Thoroughly mix 2.5 pounds Tylan 40 Sulfa-G in one ton of Type C Feed to provide 100 grams of tylosin and 100 grams of sulfamethazine per ton.

WARNING: Feeds containing Tylan 40 Sulfa-G must be withdrawn 15 days before swine are slaughtered.

Not to be used after the date printed on bag.

